
Harm Reduction & Overdose Prevention Program

Presented by: Ben Williams, Hayley King and Nathan Farley Uniting Harm Reduction & Overdose Prevention Program



Acknowledgement of Traditional Owners

We acknowledge the Traditional Owners of Country throughout Australia and recognise their continuing connection to land, waters and culture. We pay our respects to their Elders past, present and emerging.





Uniting Harm Reduction & Overdose Prevention Program

- Assertive Outreach Team initially funded under the Risk of Overdose initiative in 2018
- Provide Harm Reduction education and information (on all drug types) for the prevention of substance related harm
- Naloxone Training (clients, families and staff of AOD and other relevant services) and the provision of naloxone
- Brief interventions with clients, families, friends and staff / workers of other organisations
- Client counselling and single sessions with families
- Liaison/collaboration with other organisations providing overdose prevention services to ensure continued development of the program



Criteria / Referral pathway:


- See program flyer
- Client centred meet people where they're at
- High Risk substance use / complexity
- <u>harmreduction@vt.uniting.org</u>

Support

- Conduit to other forms of treatment / support
- More intensive client assessment, individual Treatment Planning and Case Coordination (based on CRC model) as required
- Referrals to other relevant services, treatment or harm reduction services
- Discuss treatment options (ORT, MSIR, NSP's).
- Facilitate streamlined access to AOD services



Signs and Symptoms of Opioid Overdose



- Unresponsive but awake
- Limp body, heavy nod
- Pale and /or clammy face, bluish
 - fingernails and / or lips
- Shallow or erratic breathing, or not

breathing at all

- Slow or erratic pulse / heartbeat
- Vomiting, choking sounds or

gurgling

Snoring, Loss of consciousness /

death



Naloxone... what is it?

- Naloxone (brand names Prenoxad, Nyxoid) is a drug that can temporarily reverse opioid overdose.
- Opioid antagonist Naloxone works by blocking opioids (such as heroin and oxycodone) from attaching to opioid receptors in the brain.

Naloxone reversing an overdose. Naloxone has a stronger affinity to the opioid receptors than opioids, such as heroin or oxycodone, is it knocks the opioids off the receptors for a short time (30-90 minutes). This allows the person to breathe again and reverse the overdose. naloxone opioid opioid opioid opioid receptor on brain CCPE



Naloxone..

- AMPOULES contain 400 micrograms of Naloxone
- PRENOXAD Intramuscular Injection contains 5 doses x 400 micrograms of Naloxone
- NYXOID Intranasal spray contains 2 doses x 1.8 mg of Naloxone.
- Based on its pharmacokinetics, a single dose (1.8 mg) of NYXOID is expected to deliver a similar onset for reversing an opioid overdose, that is comparable to Intramuscular Naloxone (400 micrograms).
- If administering a second dose of NYXOID it is equivalent to 5 doses of 400 micrograms of Intramuscular Naloxone.





Responding to an Opioid Overdose with NYXOID

- Check for signs of overdose
- Give 2 rescue breaths (if trained to administer CPR)
- Call an ambulance Triple 000
- Administer Nyxoid
- Place in recovery position
- Monitor and give support until the ambulance arrives
- Administer 2nd dose of Nyxoid if no improvement after
 - 2–3 minutes
- Watch for acute withdrawal symptoms











Naloxone

Take Home Naloxone program

What is it...



About the Take Home Naloxone program

The Australian Government is investing \$19.6 million over 4 years from 2022-23 to deliver the Take Home Naloxone program nationally.

From 1 July 2022:

- All community pharmacies will need to register via PPA online
- Hospital pharmacies
- Approved medical practitioners

From 1 November 2022: In Victoria naloxone will be available from the following sites

- Needle and syringe programs
- Medically Supervised Injecting Room

These services are considered Authorised Alternative Suppliers.

<u>Uniting</u>

Thank you

Contact details:

harmreduction@vt.uniting.org

ben.williams@vt.uniting.org 0466 085 557

hayley.king@vt.uniting.org 0466 920 058

nathan.farley@vt.uniting.org 0466 348 900



Resources

https://www.penington.org.au/resources/cope-overdose-first-aid/

https://www.youtube.com/watch?v=OlubISZpwQA

https://nyxoid.com.au/nyxoid-learn/

https://www.health.gov.au/initiatives-and-programs/take-homenaloxone-program/about-the-take-home-naloxone-program



